

---

**SUPPLEMENTAL LABEL**

---



**Syngenta Crop Protection, Inc.**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300

**Bravo Weather Stik®**

**Agricultural Fungicide**

Supplemental label for brassica (head and stem), cucurbits, fruiting vegetables, ginseng, horseradish, lupine and lentil, persimmon, rhubarb, and yam

*Active Ingredient:*

Chlorothalonil (tetrachloroisophthalonitrile).....	54.0%
--	-------

<i>Other Ingredients:</i>	46.0%
---------------------------	-------

Total	100.0%
-------	--------

Contains 6.0 lbs. of Chlorothalonil per gallon (720 grams per liter).

**KEEP OUT OF REACH OF CHILDREN.**

**CAUTION**

**EPA Reg. No. 50534-188-100**

**DIRECTIONS FOR USE**

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.**

**Before using Bravo Weather Stik as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application.**

CROP	DISEASE (Pathogen)	PTS PRODUCT/A (lbs a.i/A)	APPLICATION DIRECTIONS
<b>Brassica, Head and Stem</b> Broccoli Broccoli, Chinese Brussels Sprouts Cabbage Cabbage, Chinese (tight-headed varieties only) Cabbage, Chinese (napa) Cabbage, Chinese Mustard Cauliflower Cavalo Broccolo Kohlrabi	Alternaria leaf spot ( <i>Alternaria</i> spp.)  Downy mildew ( <i>Peronospora parasitica</i> )	1½ (1.125)	Use in sufficient water to obtain adequate coverage. Begin applications after transplants are set in field, or shortly after emergence of field-seeded crop, or when conditions favor disease development. Repeat at 7 to 10 day intervals (the minimum re-treatment interval is 7 days) to maintain control.  Apply by ground, air or chemigation.
	Ring spot (California only)	2 (1.5)	For field-seeded Brussels sprouts, begin applications at time of early sprout development or when conditions favor disease development. Repeat at 7 to 10 day intervals (the minimum re-treatment interval is 7 days) to maintain control.

**Specific Use Restrictions:**

- Do not apply more than 11.7 pints of Bravo Weather Stik (8.8 lbs. a.i.) per acre during each growing season.
- Do not apply within 7 days of harvest.

CROP	DISEASE (Pathogen)	PTS PRODUCT/A (lbs a.i/A)	APPLICATION DIRECTIONS
<b>Cucurbits</b> Cantaloupe Cucumber Honeydew melon Muskmelon Pumpkin Squash Watermelon Zucchini	Anthracnose ( <i>Colletotrichum</i> spp.)  Downy mildew ( <i>Pseudoperonospora cubensis</i> )  Target spot ( <i>Corynespora cassiicola</i> )	1½ to 2 (1.125 to 1.5)	Use in sufficient water to obtain adequate coverage. Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals (the minimum re-treatment interval is 7 days).  Note: Spraying mature watermelons may result in sunburn of the upper surface of the fruit. Do not apply Bravo Weather Stik to watermelons when any of the following conditions are present:
Including cultivars and/or hybrids of these.  See additional cucurbit crops below.	Alternaria leaf blight ( <i>A. cucumerina</i> )  Alternaria leaf spot ( <i>A. alternata</i> )  Cercospora leaf spot ( <i>C. citrullina</i> )  Gummy stem blight/vine decline ( <i>Didymella bryoniae</i> )  Powdery mildew ( <i>Sphaerotheca</i> only)  Scab ( <i>Cladosporium cucumerinum</i> )	2 to 3 (1.5 to 2.25)	<ol style="list-style-type: none"> <li>1. Intense heat and sunlight</li> <li>2. Drought conditions</li> <li>3. Poor vine canopy</li> <li>4. Other crop and environmental conditions which may be conducive to increased natural sunburn</li> </ol> Do not combine Bravo Weather Stik with anything except water for application to watermelons unless your prior use has shown the combination to be non-injurious to watermelons under your conditions of use.  Apply by ground, air or chemigation.

**Additional cucurbit crops:** Chayote, Chinese waxgourd, Gourds, *Momordica* spp. (Bitter melon, Balsam apple)

**Specific Use Restrictions:**

- Do not apply more than 21 pints of Bravo Weather Stik (15.75 lbs. a.i.) per acre during each growing season.
- Bravo Weather Stik may be applied the day of harvest (0 day PHI).

CROP	DISEASE (Pathogen)	PTS PRODUCT/A (lbs a.i/A)	APPLICATION DIRECTIONS
<b>Fruiting Vegetables:</b> (except tomato) Eggplant Groundcherry Okra Pepino Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper) Tomatillo	Anthracnose ( <i>Colletotrichum</i> spp.)  Botrytis leaf mold ( <i>Botrytis cinerea</i> )  Cercospora leaf spot ( <i>Cercospora</i> spp.)  Powdery mildew ( <i>Leveillula taurica</i> )	1½ (1.125)	Use in sufficient water to obtain adequate coverage. Begin applications as a foliage, flower, and fruit spray when disease is expected. Repeat applications at 7-10 day intervals.  Apply by ground, air or chemigation.

**Specific Use Restrictions:**

- Do not apply more than 12 pints of Bravo Weather Stik (9.0 lbs. a.i.) per acre during each growing season.
- Do not apply within 3 days of harvest (3 day PHI).

<b>Ginseng</b>	Alternaria blight ( <i>Alternaria panax</i> )  Gray mold ( <i>Botrytis cinerea</i> )	2 (1.5)	Use in sufficient water to obtain adequate coverage. Begin applications when disease first threatens, and repeat at 7-10 day intervals as disease pressure warrants.
----------------	--	------------	--

**Specific Use Restrictions:**

- Do not apply more than 16 pints of Bravo Weather Stik (12 lbs. a.i.) per acre during each growing season.
- Do not apply within 14 days of harvest (14 day PHI).

<b>Horseradish</b>	Ramularia stem and leaf spot ( <i>Ramularia armoraciae</i> )	3 (2.25)	Use in sufficient water to obtain adequate coverage. Begin applications when disease first threatens, and repeat at 7-10 day intervals as disease pressure warrants.
--------------------	---	-------------	--

**Specific Use Restrictions:**

- Do not apply more than 24 pints of Bravo Weather Stik (18 lbs. a.i.) per acre during each growing season.
- Do not apply within 14 days of harvest (14 day PHI).

<b>Lupine and Lentil</b>	Anthracnose ( <i>Colletotrichum gloeosporioides</i> )  Ascochyta ( <i>Ascochyta pisi</i> )	1 to 1.5 (0.75 to 1.125)	Use in sufficient water to obtain adequate coverage. Begin applications when disease first threatens, and repeat at 7-10 day intervals as disease pressure warrants.
--------------------------	--	-----------------------------	--

**Specific Use Restrictions:**

- Do not apply more than 8 pints of Bravo Weather Stik (6 lbs. a.i.) per acre during each growing season.
- Do not apply within 14 days of harvest (14 day PHI).

CROP	DISEASE (Pathogen)	PTS PRODUCT/A (lbs a.i/A)	APPLICATION DIRECTIONS
Persimmon	Cercospora leaf spot ( <i>Cercospora fuliginosa</i> )	1.25 (0.94)	Use in sufficient water to obtain adequate coverage. Begin applications when disease first threatens, and repeat at 14 day intervals as disease pressure warrants.

**Specific Use Restrictions:**

- Do not apply more than 6.25 pints of Bravo Weather Stik (4.7 lbs. a.i.) per acre during each growing season.
- Do not apply within 14 days of harvest (14 day PHI).
- May be applied to persimmon only in the states of Florida and Hawaii.
- Aerial applications require the use of a minimum of 10 gallons per acre.

Rhubarb	Ramularia leaf spot ( <i>Ramularia rhei</i> )  Ascochyta ( <i>Ascochyta rhei</i> )	3 (2.25)	Use in sufficient water to obtain adequate coverage. Begin applications when disease first threatens, and repeat at 7-10 day intervals as disease pressure warrants.
---------	--	-------------	--

**Specific Use Restrictions:**

- Do not apply more than 18 pints of Bravo Weather Stik (13.5 lbs. a.i.) per acre during each growing season.
- Do not apply within 30 days of harvest (30 day PHI).

Yam	Anthrachnose ( <i>Colletotrichum gloeosporioides</i> )	1 to 1.25 (0.75 to 1.125)	Use in sufficient water to obtain adequate coverage. Begin applications when disease first threatens, and repeat at 10-14 day intervals as disease pressure warrants.
-----	---	------------------------------	---

**Specific Use Restrictions:**

- Do not apply more than 15 pints of Bravo Weather Stik (11.25 lbs. a.i.) per acre during each growing season.
- Do not apply within 7 days of harvest (7 day PHI).

Bravo Weather Stik® and the Syngenta logo are trademarks of a Syngenta Group Company

©2009 Syngenta

Syngenta Crop Protection, Inc.  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300

SCP 50534-188A-S5 0109

BRA WEA 50534-188A-S5-0109 – df – 3/9/09